

medium. This model predicts how the SBS Stokes intensity depend upon the laser pump intensity and upon the physical properties of the SBS medium. A brief study of a numerical approximation to the Brillouin wave equations was presented as MATLAB code and simulation result.

Conflict of Interest

The author declares no conflict of interest.

References

- [1] Debye, P., & Sears, F. W. (1932). On the scattering of light by supersonic waves. *Proceedings of the Nat. Aced. Sci.*, 409414.
- [2] McCurdy, A. H. (2005). Modeling of Spontaneous Brillouin scattering in optical fibers with arbitrary radial index profile. *Journal of Lightwave Technology*, 23(11).
- [3] Nieves, O. A., Arnold, M. D., Steel, M. J., Schmidt, M. K., & Poulton, C. G. (2020). Noise and pulse dynamics in backward spontaneous Brillouin scattering. Retrieved from <https://www.researchgate.net/publication/345989265>
- [4] Bongrond, I., Picholle, É., & Montes, C. (2002). Coupled longitudinal and transverse stimulated Brillouin scattering in single-mode optical fibers. *The European Physical Journal D - Atomic, Molecular, Optical and Plasma Physics*, 20, 121–127.
- [5] Tang, C. L. (1966). Saturation and spectral characteristics of the stokes emission in the stimulated Brillouin process. *Journal of Applied Physics*, 37, 2945.
- [6] Baldwin, G. C. (1969). *An Introduction to Nonlinear Optics*. New York: Plenum Press.
- [7] Boyd, R. W. (2003). *Nonlinear Optics* (2nd ed.). San Diego: Academic Press.

Copyright © 2023 by the authors. This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).